



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Placing  $\tan \theta = t$ ,  $d\theta = \frac{dt}{1+t^2}$ ,  $\sin \theta = \frac{t}{(1+t^2)^{1/2}}$ , and  $\cos \theta = \frac{1}{(1+t^2)^{1/2}}$ .

Integrating, we obtain,

$$y+c = -\frac{2}{3} [(\tan \frac{1}{2} \theta)^{-\frac{1}{3}} + (\tan \frac{1}{2} \theta)^{\frac{1}{3}}] \frac{2 \cdot 16^{\frac{1}{6}} a^2}{27 \cdot 2^{\frac{1}{3}}} = -\frac{a^2}{36} \left[ \left( \tan \frac{1}{2} \theta \right)^{-\frac{1}{3}} + \left( \tan \frac{1}{2} \theta \right)^{\frac{1}{3}} \right]$$

from which, by substitution, the solution may be completed.

III. Solution by WILLIAM HOOVER, Athens, Ohio.

Let  $dy/dx = p$ , as usual; then  $p^3 - axp + x^3 = 0$ .

The condition for equal values of  $p$  from this cubic is

$$\frac{p^6}{4} = \frac{a^3 p^3}{27}, \text{ or } p = \frac{dy}{dx} = \frac{a}{3} \sqrt[3]{4}.$$

Integrating,  $\frac{3y}{a\sqrt[3]{4}} = x+c$ , or,  $27y^3 = 4a^3(x+c)^3$ .

Also solved by S. G. Barton.

---

## MECHANICS.

---

255. Proposed by the late G. B. ZERR, Ph. D.

Assuming the resilience of volume of mercury to be constant at all depths and to be  $54.20 \times 10^{10}$  in C. G. S. units and that a mile = 160933 centimeters. Find the depth of an ocean of mercury at a point where its density is double the surface density, 13.596.

Solution by B. F. FINKEL, Ph. D., Drury College.

Let  $x_0$  be the length of a column of mercury of one square centimeter cross-section and uncompressed, such that its weight is sufficient to compress a cubic centimeter to half its volume, which is equivalent to doubling its density.

By Hooke's law,  $p = E \frac{\Delta v}{v} = \frac{1}{2} E$ , where  $E$  is the coefficient of elasticity. But  $p = 13.596 g x_0$ .

$\therefore x_0 = \frac{E}{27.192g}$ . Let  $\Delta x_0$  = the amount of compression of the column by virtue of its own weight, and let  $x$  = the length of any portion of the column measured from the surface downward, and let  $dx$  be an element of length of this column. Then the strain,  $s$ , in this element is

$s = \frac{d(\Delta x_0)}{dx}$ . But by Hooke's Law,  $s = \frac{p}{E}$ . Hence,  $\frac{d(\Delta x_0)}{dx} = \frac{p}{E}$

$= \frac{13.596gx}{E}$ , or  $d(\Delta x_0) = \frac{13.596gxdx}{E}$ , and

$$\therefore \Delta x_0 = \frac{13.596g}{E} \int_0^{x_0} x dx = \frac{13.596g(\frac{1}{2}x_0^2)}{E} = \frac{6.798gx_0^2}{E}$$

$\therefore$  The depth of the ocean required is  $x = x_0 - \Delta x_0$

$$\begin{aligned} &= \frac{E}{27.192g} - \frac{6.798g}{E} \left( \frac{E}{27.192g} \right)^2 = \frac{E}{27.192g} \left[ 1 - \frac{1}{4} \right] \\ &= \frac{3E}{108.768g} \text{ cm.} = \frac{3E}{108.768 \times 160933 \times g} \text{ mi.} \end{aligned}$$

Using  $g=981$ , and  $E=33 \times 10^{10}$ , we have  $x=61.8$  miles.

REMARK. We are unable to get data from which to obtain Dr. Zerr's value for the volume resilience. The value of  $E$  as given in Kimball's *Physics*, page 158, and which we have used in the above solution, is very different from what would be obtained from Dr. Zerr's value for volume resilience.

---

## NOTES AND NEWS.

---

The fifth regular meeting of the Southwestern Section of the American Mathematical Society was held at Washington University, St. Louis, Missouri, on Saturday, December 2, 1911. S.

The eighteenth summer meeting of the American Mathematical Society was held at Poughkeepsie, New York, on September 12-13, 1911. There were about thirty-five members in attendance and twenty-six papers were presented. S.

The twenty-ninth regular meeting of the Chicago Section of the American Mathematical Society will be held at the University of Chicago on Friday and Saturday, December 29, 30, 1911. Titles and abstracts of papers to be presented at this meeting should be sent to the secretary of the Section, Professor H. E. Slaught, the University of Chicago, Chicago, Illinois. S.